



Phylogenetic Analysis and Polyphasic Characterization of *Clavibacter michiganensis* Strains Isolated from Tomato Seeds Reveal that Nonpathogenic Strains Are Distinct from *C. michiganensis* subsp. *michiganensis*

Submitted by Emmanuel Lemoine on Thu, 02/12/2015 - 13:08

Titre	Phylogenetic Analysis and Polyphasic Characterization of <i>Clavibacter michiganensis</i> Strains Isolated from Tomato Seeds Reveal that Nonpathogenic Strains Are Distinct from <i>C. michiganensis</i> subsp. <i>michiganensis</i>
Type de publication	Article de revue
Auteur	Jacques, Marie-Agnès [1], Durand, Karine [2], Orgeur, Geoffrey [3], Balidas, Samuel [4], Fricot, Céline [5], Bonneau, Sophie [6], Quillévéré, Anne [7], Audusseau, Corinne [8], Olivier, Valérie [9], Grimault, Valérie [10], Mathis, René [11]
Editeur	American Society for Microbiology
Type	Article scientifique dans une revue à comité de lecture
Année	2012
Langue	Anglais
Date	2012/01/12
Numéro	23
Pagination	8388 - 8402
Volume	78
Titre de la revue	Applied and Environmental Microbiology
ISSN	0099-2240

Résumé en anglais	<p>The genus <i>Clavibacter</i> comprises one species and five subspecies of plant-pathogenic bacteria, four of which are classified as quarantine organisms due to the high economic threat they pose. <i>Clavibacter michiganensis</i> subsp. <i>michiganensis</i> is one of the most important pathogens of tomato, but the recommended diagnostic tools are not satisfactory due to false-negative and/or -positive results. To provide a robust analysis of the genetic relatedness among a worldwide collection of <i>C. michiganensis</i> subsp. <i>michiganensis</i> strains, relatives (strains from the four other <i>C. michiganensis</i> subspecies), and nonpathogenic <i>Clavibacter</i>-like strains isolated from tomato, we performed multilocus sequence-based analysis and typing (MLSA and MLST) based on six housekeeping genes (<i>atpD</i>, <i>dnaK</i>, <i>gyrB</i>, <i>ppK</i>, <i>recA</i>, and <i>rpoB</i>). We compared this “framework” with phenotypic and genotypic characteristics such as pathogenicity on tomato, reaction to two antisera by immunofluorescence and to five PCR identification tests, and the presence of four genes encoding the main <i>C. michiganensis</i> subsp. <i>michiganensis</i> pathogenicity determinants. We showed that <i>C. michiganensis</i> subsp. <i>michiganensis</i> is monophyletic and is distinct from its closest taxonomic neighbors. The nonpathogenic <i>Clavibacter</i>-like strains were identified as <i>C. michiganensis</i> using 16S rRNA gene sequencing. These strains, while cross-reacting with <i>C. michiganensis</i> subsp. <i>michiganensis</i> identification tools, are phylogenetically distinct from the pathogenic strains but belong to the <i>C. michiganensis</i> clade. <i>C. michiganensis</i> subsp. <i>michiganensis</i> clonal complexes linked strains from highly diverse geographical origins and also strains isolated over long periods of time in the same location. This illustrates the importance of seed transmission in the worldwide dispersion of this pathogen and its survival and adaptation abilities in a new environment once introduced.</p>
URL de la notice	http://okina.univ-angers.fr/publications/ua7824 [12]
DOI	10.1128/AEM.02158-12 [13]
Lien vers le document	http://dx.doi.org/10.1128/AEM.02158-12 [13]

Liens

- [1] <http://okina.univ-angers.fr/m.jacques/publications>
- [2] [http://okina.univ-angers.fr/publications?f\[author\]=11808](http://okina.univ-angers.fr/publications?f[author]=11808)
- [3] [http://okina.univ-angers.fr/publications?f\[author\]=12398](http://okina.univ-angers.fr/publications?f[author]=12398)
- [4] [http://okina.univ-angers.fr/publications?f\[author\]=12399](http://okina.univ-angers.fr/publications?f[author]=12399)
- [5] [http://okina.univ-angers.fr/publications?f\[author\]=12400](http://okina.univ-angers.fr/publications?f[author]=12400)
- [6] [http://okina.univ-angers.fr/publications?f\[author\]=12368](http://okina.univ-angers.fr/publications?f[author]=12368)
- [7] [http://okina.univ-angers.fr/publications?f\[author\]=12401](http://okina.univ-angers.fr/publications?f[author]=12401)
- [8] [http://okina.univ-angers.fr/publications?f\[author\]=12402](http://okina.univ-angers.fr/publications?f[author]=12402)
- [9] [http://okina.univ-angers.fr/publications?f\[author\]=12403](http://okina.univ-angers.fr/publications?f[author]=12403)
- [10] [http://okina.univ-angers.fr/publications?f\[author\]=12404](http://okina.univ-angers.fr/publications?f[author]=12404)
- [11] [http://okina.univ-angers.fr/publications?f\[author\]=12405](http://okina.univ-angers.fr/publications?f[author]=12405)
- [12] <http://okina.univ-angers.fr/publications/ua7824>
- [13] <http://dx.doi.org/10.1128/AEM.02158-12>

Publié sur *Okina* (<http://okina.univ-angers.fr>)